

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 January 2002 (31.01.2002)

PCT

(10) International Publication Number
WO 02/08434 A2

- (51) International Patent Classification⁷: C12N 15/86 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (21) International Application Number: PCT/EP01/08124
- (22) International Filing Date: 13 July 2001 (13.07.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 00115626.4 20 July 2000 (20.07.2000) EP
- (71) Applicant: ARTEMIS PHARMACEUTICALS GMBH [DE/DE]; Neurather Ring 1, 51063 Köln (DE).
- (72) Inventors: HOBOM, Gerd; Unterfeldstrasse 13, 79106 Freiburg (DE). MENKE, Anette; Schillingstrasse 37-39, 50670 Köln (DE). MEYER-ROGGE, Sabine; Oberpforte 7, 35321 Laubach-Münster (DE).
- (74) Agents: HELBING, Jörg et al.; Von Kreisler Selting Werner, Postfach 10 22 41, 50462 Köln (DE).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/08434 A2

(54) Title: RECOMBINANT INFLUENZA VIRUSES WITH BICISTRONIC vRNAs CODING FOR TWO GENES IN TANDEM ARRANGEMENT

(57) Abstract: The invention relates to recombinant influenza viruses for high-yield expression of incorporated foreign gene(s), which are genetically stable in the absence of any helper virus and which comprise at least one viral RNA segment being a tandem bicistronic RNA molecule coding for two genes in tandem, in said tandem bicistronic RNA molecule one of the standard viral genes being in covalent junction with a foreign, recombinant gene and having an upstream splice donor and a downstream splice acceptor signal surrounding the proximal coding region. The invention further provides a method for obtaining attenuated viruses which resist reassortment dependent progeny production in case of chance superinfections by wild-type influenza viruses; a method for the production of said recombinant influenza viruses; pharmaceutical compositions comprising said recombinant influenza viruses; and the use of said recombinant influenza viruses for preparing medicaments for vaccination purposes.

Recombinant Influenza Viruses with Bicistronic vRNAs Coding for Two Genes in Tandem Arrangement

5

Field of the Invention

The invention relates to recombinant influenza viruses for high-yield expression of incorporated foreign gene(s), which are genetically stable in the absence of any helper virus and which comprise at least one viral RNA segment being a tandem bicistronic RNA molecule coding for two genes in tandem, in said tandem bicistronic RNA molecule one of the standard viral genes being in covalent junction with a foreign, recombinant gene and having an upstream splice donor and a downstream splice acceptor signal surrounding the proximal coding region. In particular the above tandem bicistronic RNA molecule contains one of the standard viral genes in distal mRNA position behind a foreign, recombinant gene in proximal position, or vice versa, both in antisense orientation with regard to the viral RNA within the virus. For simultaneous expression of both genes the proximal reading frame is flanked by splice donor and acceptor signals which have the quality to allow a partial yield of spliced mRNA only, i.e., resulting in the presence of both, spliced and unspliced mRNA simultaneously.

The invention further provides a method for obtaining attenuated viruses which resist reassortment dependent progeny production in case of chance superinfections by wild-type influenza viruses; a method for the production of said recombinant influenza viruses; pharmaceutical compositions comprising said recombinant influenza viruses; and the use of said recombinant influenza viruses for preparing medicaments for vaccination purposes.

30

Technical Background

Redesigning influenza virus into a vector system for expression of foreign genes similar to what has been achieved in several other thoroughly studied viruses such as adenovirus, retrovirus, Semliki Forest virus or Rabies virus has the advantage of an industrially well established mode of cheap propagation for influenza in fertilized chicken eggs leading to rather high titers (above 10^{10} /ml). On the other hand none of the constituent vRNA segments may be deleted from the influenza genome according to our present knowledge, and give room for large-size foreign insertions.

Only small fragments of foreign polypeptide chains such as B cell epitopes (10 to 15 amino acids) may be inserted into selected positions within two of the viral proteins, i.e. in exchange for one of the variable antigenic regions located at the surface of hemagglutinin (Muster et al., Mucosal model of immunization against human immunodeficiency virus type 1 with a chimeric influenza virus, J. Virol. 69 (11), 6678-6686 (1995)), or into the stalk sequence of viral neuraminidase (Garcia-Sastre and Palese, The cytoplasmic tail of the neuraminidase protein of influenza A virus does not play an important role in the packaging of this protein into viral envelopes, Virus Res. 37, 37-47 (1995)), and be stably maintained as functional fusion proteins. Constructs of this kind turned out to be useful for experimental vaccination in a few cases studied, but only rather few clearly defined epitope sequences (of ten to twelve amino acids each) are known today, and some of them might also be misfolded within such restricted fusion protein positions, or in other cases interfere with formation of the correct tertiary structure and function of their host polypeptide chains.

Incorporation of a full-size foreign protein into influenza virus via reverse genetics, encoded by an independent ninth vRNA molecule in addition to its regular set of eight standard vRNA segments is without special provisions only transiently possible (Luytjes et al., Amplification, expression, and packaging of a foreign gene by influenza virus. Cell 59, 1107-1113 (1989); Enami et al., An influenza virus containing nine

different RNA segments, Virology 185, 291-298 (1991)). In the absence of a continuous selective pressure any additional recombinant vRNA segment cannot be stably maintained as long as the wildtype promoter sequence is used on that ninth vRNA segment, and it will inadvertently be lost after
5 few steps of viral propagation.

Using a different system of influenza reverse genetics developed in our laboratory (Zobel et al., RNA polymerase I catalysed transcription of insert viral cDNA, Nucleic Acids Res. 21, 3607-3614 (1993); Neumann et al., RNA polymerase I-mediated expression of influenza viral RNA molecules,
10 Virology 202, 477-479 (1994)), which was built around *in vivo* synthesis of recombinant vRNA molecules by cellular RNA polymerase I transcription of the respective template cDNA constructs, modified terminal viral RNA sequences (hereinafter "promoter-up mutations" or promoter-up variants") have been designed by nucleotide substitutions (Neumann and
15 Hobom, Mutational analysis of influenza virus promoter elements *in vivo*, J. Gen. Virol. 76, 1709-1717 (1995); WO 96/10641). The above promoter-up variants carry up to five nucleotide substitutions (in promoter-up variant 1920; see Flick and Hobom, J. Gen. Virol. 80, 2565-2572 (1999)). When these promoter-up variants are attached to a
20 recombinant ninth vRNA segment its increased transcription and amplification rate will not only compensate for the losses suffered spontaneously, but even cause accumulation of the foreign vRNA segment during simple viral passaging, in the absence of any selection.

25 However, due to its over-replication relative to all of the regular influenza vRNA segments (which of course are connected to wild-type promoter sequences) after catching up with the others the foreign segment will become over-abundant. This increasingly will result in viral particles that have incorporated several copies of recombinant vRNA, but no longer have
30 a full set of all eight standard segments incorporated among an average of about 12-15 vRNA molecules present within a virion. Such particles are defective and will not result in plaque formation, hence after an initial

increase of recombinant viral particles during the first steps of propagation a dramatic decrease is observed, usually at the third or fourth step of viral passaging, depending on the size of the recombinant vRNA and the level of the promoter-up mutation attached.

5

A balanced situation with regard to the insert length and the level of promoter activity can be achieved, and has been propagated in a particular case over 11 passages, with essentially stable levels of recombinant viruses among a majority of helper viruses (around 80%) during these steps. If a full-level promoter-up mutation is used (1104 or the variant 1920, see below) a balanced-level propagation is reached in conjunction with a recombinant vRNA size of 4000 nucleotides (Maysa Azzeh, Ph.D. Thesis, Univ. Giessen (2000)).

In all of these preparations, both in transiently achieved increased yields (up to 40% of recombinants after three or four steps of viral passage), and in a balanced propagation of recombinant influenza viruses (10 - 20%) the respective viral progeny inadvertently constitute mixtures with a majority of non-recombinant helper viruses. These result both from a statistical mode of packaging vRNA molecules into a virion (the ninth segment may not be co-packaged), and from the fraction of cells solely infected by helper virus.

To solve the problems of fractional yields and of instability during viral propagation of recombinant influenza, it was suggested to use a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus and which comprises at least one viral RNA segment being an ambisense RNA molecule (designated "ambisense RNA segment") and containing one of the standard viral genes in sense orientation and a foreign, recombinant gene in anti-sense orientation, or *vice versa*, in overall convergent arrangement (PCT/EP00/01903). The ambisense RNA segment preferably should contain the promoter-up mutations. The PCT/EP00/01903 moreover discloses a method of constructing specific

influenza carrier (helper) strains carrying one or more ribozyme target sites (of type one) in vRNA flanking positions comprising

- (a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, carrying two different 3' promoter sequences in tandem (an external promoter-up variant and an internal wild-type promoter), which are separated by a second type of ribozyme target sequence, and which carry the said internal ribozyme target sites of type one;
- (b) followed by infection of an influenza wildtype strain;
- (c) thereafter amplification through simple steps of viral propagation; and
- 10 (d) finally isolation through removal of their external 3' promoter sequence by ribozyme cleavage through infection of cells expressing ribozyme type 2, followed by plaque purification.

The resulting special helper virus strains carrying a vRNA segment with external ribozyme target sites of type 1 in exchange for the equivalent regular vRNA molecule are then used for the rescue of ambisense RNA molecules. These are exclusively maintained in the recombinant viruses after passage of viral propagation through ribozyme (type 1) containing host cells, which will destroy the sensitive vRNA molecules of the specially prepared helper viruses.

However, the above ambisense constructs are susceptible to (intra-nuclear) mRNA double-strand formation, which will partially reduce the expression rates of both the ambisense genes, in particular the gene driven by the (weaker) cRNA promoter. The fluctuating extent of this effect made it difficult to bring the expression rate of the influenza gene within the ambisense segment into balance with other influenza genes. This was the problem to be solved with the present invention.

30 **Summary of the Invention**

Starting out from two observations in this laboratory which are discussed above and which concern two hitherto unsuspected properties of influenza

viral RNA polymerase in its interaction with terminally adapted influenza-specific RNA molecules, stable recombinant influenza viruses were found, which solve the above problems.

- 5 The recombinant viruses of the present invention can be used for cheap propagation in fertilized eggs, either for production of those recombinant viruses themselves or for production of foreign proteins or glycoproteins encoded by them, and hence find application in (glyco)protein production or in providing vector systems for somatic gene therapy or in being used
10 as vaccination agents.

Thus, the present invention provides

- (1) a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper
15 virus and which comprises at least one viral RNA segment being a bicistronic RNA molecule coding for two genes in tandem arrangement (hereinafter "tandem bicistronic RNA segment" or "tandem RNA segment"), in said tandem RNA segment one of the standard viral genes being in covalent junction with a foreign, recombinant gene and said
20 tandem RNA segment having an upstream splice donor and a downstream splice acceptor signal surrounding the proximal coding region;
- (2) a preferred embodiment of the recombinant influenza virus defined in (1) above, in which the terminal viral RNA sequences of said at least one tandem RNA segment, which are active as the promoter signal, have been
25 modified by nucleotide substitutions in up to five positions, resulting in improved transcription rates of both the vRNA promoter as well as the cRNA promoter as present in the complementary sequence;
- (3) a method for the production of recombinant influenza viruses as defined in (1) and (2) above comprising
- 30 (a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, in antisense, or in sense tandem design,

- (b) followed by infection with an influenza carrier strain constructed to include flanking ribozyme target sequences in the corresponding viral RNA segment, i.e., coding for the same viral gene as present in the tandem segment distal position, and
- 5 (c) thereafter selective vRNA inactivation through ribozyme cleavage;
- (4) a pharmaceutical composition comprising a recombinant influenza virus as defined in (1) and (2) above;
- (5) the use of a recombinant influenza virus as defined in (1) and (2) above for preparing a medicament for vaccination purposes;
- 10 (6) the use of a recombinant influenza virus as defined in (1) and (2) above for preparing agents for somatic gene therapy;
- (7) the use of a recombinant influenza virus as defined in (1) and (2) above for preparing agents for transfer and expression of foreign genes into cells (abortively) infected by such viruses;
- 15 (8) the use of a recombinant influenza virus as defined in (1) and (2) above for preparing agents for transfer and expression of RNA molecules into cells infected by such viruses;
- (9) a method for the production of proteins or glycoproteins which comprises utilizing a recombinant influenza virus as defined in (1) and (2)
- 20 above as expression vector;
- (10) a method for preventing and/or treating influenza which comprises administering a recombinant influenza virus as defined in (1) and (2) above to the mammal to be treated, i.e., a vaccination method utilizing said recombinant virus;
- 25 (11) a method for somatic gene therapy, which method comprises subjecting the organism to be treated with a recombinant influenza virus as defined in (1) and (2) above;
- (12) a method for transfer and expression of foreign genes into cells, and for transfer and expression of RNA molecules into cells, which method
- 30 comprises infecting the cells with a recombinant influenza virus as defined in (1) and (2) above;

- (13) use of a recombinant influenza virus as defined in (1) and (2) above for preparing agents for autologous immunotherapy;
- (14) a method for an immunotherapy which comprises *ex vivo* infection of immune cells with a recombinant influenza virus as defined in (1) and (2) above; and introduction of the transduced cells into the patient; and
- (15) a method for the induction of antibodies which comprises utilizing a recombinant influenza virus as defined in (1) and (2) above as an immunogen.
- 10 The invention is described in more detail below.

Brief Description of the Figures

Fig. 1 shows the basepair substitution analysis according to the vRNA 'corkscrew' structure:

(A) 'Corkscrew' conformation of the vRNA promoter drawn against a schematic indication of interacting tripartite viral polymerase. Paired positions exchanged in individual experiments are indicated by numbers, nucleotides $\bar{3}$ or $\bar{8}$ are counted from the 3' end. pHL2024 containing promoter-up mutation '1104' is used as the reference construct (=100%) in all of the CAT assays, while pHL2428 represents the wild-type promoter structure.

(B) CAT analysis of a series of substitution variants in positions 3 and 8 from the 5' end as indicated above the lanes; 50 μ l of cell lysate obtained from 10^6 MDCK cells infected in the first viral passage with recombinant viral progeny.

(C) pHL2024 and pHL1920 comparative CAT analysis, in 100 fold dilution relative to (B), i.e., obtained from 0.5 μ l of cell lysate in 3 h reaction time.

Fig. 2: Vector plasmid pHL1920, the exact sequence of the 3888 bps circular DNA is shown in SEQ ID NO: 20

Fig. 3 shows the genetic structure and the RNA transcription products of influenza model tandem expression constructs. Heavy lines for the plasmid cDNA constructs refer to double-stranded DNA, while single-stranded RNA molecules are represented by thin lines, and their 5' to 3' directionalities are marked by arrows. Standard modifications at their 5' and 3' ends are indicated by a dot (5' cap structure) and A_n (3' polyadenylation), both are absent in the primary anti-sense transcription product, the viral RNA (vRNA), which is transcribed by cellular RNA polymerase I (RPoI). Full-length mRNA₁ is synthesized by influenza viral polymerase (virPo), and a partial splice reaction results in a functional yield of shorter mRNA₂ molecules. While both of the reporter genes are indicated on the DNA level, together with the positions of splice donor (D) and acceptor (A) signal sequences as well as the promoter (p_i) and terminator (t_i) elements for RNA polymerase I start and stop, on the RNA level only those genes and splice signals are marked that are actually translated into protein or actively involved in splicing. The chloramphenicol-acetyltransferase gene (CAT) has been inserted in proximal position in pHL3196 and pHL3235, and in distal position in pHL3224 and pHL3236 (see Figs. 5 to 8), while green fluorescent protein (GFP) in each case is located in alternate location. All vRNA molecules - and hence, also the cDNA constructs - carry sequence variations at their 3' ends, which together constitute the 1104 promoter-up mutations: G3A, U5C, C8U (nucleotide positions counted from the 3' vRNA end). pHL3235 and pHL3236 vRNAs are extended in size by about 1000 nucleotides of untranslated sequence relative to pHL3196 and pHL3226: 2600 instead of 1600 nucleotides in lengths. For full-size representation of circular plasmid DNAs see Figs. 5-8, for CAT expression data of all infected by recombinant influenza viruses carrying the respective viral RNAs see Fig 4.

Fig. 4 shows the CAT assay results for the group of tandem vRNA plasmid constructs as described in the Example. In particular, the ratio between chloramphenicol (bottom line) and acetylchloramphenicol (upper three

lines) in a flash-CAT assay, after the 2nd (A) and 4th (B) passage of recombinant viruses carrying the reportergene CAT, can be determined from said figure. The following constructs were utilized:

5 pHL1844 (control): monocistronic CAT-construct downstream of promoter variant 1104.

pHL3196: tandem construct, p-CAT-GFP resulting in a vRNA having a total length of 1530 nucleotides (not "extended"), see also Fig. 5.

pHL3235: tandem construct, p-CAT-GFP resulting in a vRNA having a total length of 2550 nucleotides ("extended"), see also Fig. 7.

10 pHL3224: tandem construct, p-CAT-GFP resulting in a vRNA having a total length of 1700 nucleotides (not "extended"), see also Fig. 6.

pHL3236: tandem construct, p-CAT-GFP resulting in a vRNA having a total length of 2720 nucleotides ("extended"), see also Fig. 8.

15 pHL2899: ambisense construct, p_v-CAT→ ←GFP-p_c resulting in an RNA having a total length of 1500 nucleotides.

pHL2960: ambisense construct, p_v-CAT→ ←GFP-p_c resulting in an RNA having a total length of 1500 nucleotides.

The five constructs on the left side were transfected into the cell DNA without the use of "booster" plasmides, the four constructs on the right side were, however, transfected with the "booster" plasmides, which gives
20 a jump-start of the constructs due to recombinant vRNA amplification prior to helper virus injection, equivalent to an advantage of about two passages. The "booster" plasmides comprise expression constructs for the nucleoprotein as well as the three subunits of influenza viral polymerase, each downstream of an RNA polymerase II promoter and in an mRNA
25 forming cassette.

While the ambisense construct having the CAT-reporter gene in the weaker position, i.e. behind the cRNA promoter (pHL2899), is only expressed moderately, this is not the case in the respective tandem
30 construct having the CAT-reporter gene in the weaker position, viz. pHL3224 or pHL3236. Further, the "extension" of the vRNA by 1020 non-

translated nucleotides (at the 3' end) is tolerated without significant decrease of expression (see pHL3235 versus pHL3196).

Fig. 5: Vector plasmid pHL3196, the exact sequence of the 4500 bps
5 circular DNA is shown in SEQ ID NO:21.

Fig. 6: Vector plasmid pHL3224, the exact sequence of the 4721 bps
circular DNA is shown in SEQ ID NO:22.

10 Fig. 7: Vector plasmid pHL3235, the exact sequence of the 5517 bps
circular DNA is shown in SEQ ID NO:23.

Fig. 8: Vector plasmid pHL3236, the exact sequence of the 5699 bps
circular DNA is shown in SEQ ID NO:24.

15

Detailed Description of the Invention

According to the present invention "influenza virus" embraces influenza A virus, influenza B virus and influenza C virus, with influenza A virus being preferred.

20

"Bicistronic" according to the present invention refers to a viral RNA segment, vRNA, cRNA or mRNA that includes two independent genes in covalent junction; in a preferred version one of these genes is of viral origin, while the other one codes for a foreign, recombinant gene product.

25

"Proximal" and "proximal position" according to the present invention refers to the 5' portion of one of the genes in the bicistronic viral mRNA, i.e., ahead (upstream) of the second gene in "distal position".

30 A "mammal" according to the present invention includes humans and animals. "Organism" embraces prokaryotic and eukaryotic systems as well

as multicellular systems such as vertebrates (including mammals) and invertebrates.

"Infected cells" and "infecting cells" according to the present invention
5 also include "abortively infected cells" and "abortively infecting cells", respectively.

In a preferred influenza virus according to embodiment (1) at least one of the regular viral RNA segments is replaced by a tandem RNA segment
10 which contains one of the standard viral genes in distal position, and a foreign, recombinant gene in proximal position, both in anti-sense orientation, or vice-versa. It is moreover preferred that in the tandem RNA molecule said foreign recombinant gene is covalently bound to one of the viral genes while the original vRNA segment coding for the same gene is
15 deleted from the recombinant virus by specific ribozyme cleavage.

The foreign gene(s) in tandem covalent junction with the viral gene(s) preferably code for proteins and/or glycoproteins which are secreted from cells infected with the recombinant virus, such as lymphokines, or code for
20 glycoproteins that are incorporated into the virion as well as the plasma membrane of the infected cell. In another preferred embodiment the foreign gene(s) in tandem covalent junction with the viral gene(s) code for proteins or artificial polypeptides designed to support an efficient presentation of inherent epitopes at the surface of infected cells, for
25 stimulation of B cell and/or T cell response. Such proteins or artificial polypeptides constitute for instance a tumor antigen or an artificial oligomeric series of T cell epitopes that have been identified within a polypeptide chain. Finally, the foreign gene(s) may be suitable for transfer and expression of RNA molecules, including antisense RNAs and
30 ribozymes, into cells. Such recombinant influenza viruses are suitable for sequence specific gene silencing, for example by antisense or RNA interference mechanisms.

A preferred recombinant virus of the invention is where in the regular viral RNA segments one or both of the standard glycoproteins hemagglutinin and neuraminidase have been exchanged, preferably into fusion
5 glycoproteins consisting of an anchor segment derived from hemagglutinin and an ectodomain obtained from the foreign source, viral or cellular, or in which such recombinant glycoprotein has been inserted as a third molecular species in addition to the remaining standard components.

10 As set forth in embodiment (2) above, a preferred recombinant virus of the invention is where the terminal viral RNA sequences, which are active as promoter signal, have been modified by nucleotide substitution in up to 5 positions, resulting in improved transcription rates (of both the vRNA promoter and in the cRNA promoter as present in the complementary
15 sequence) as well as enhanced replication and/or expression rates relative to the wild-type sequence. Said modified terminal viral RNA sequences differ from the wild-type sequence in that in said tandem vRNA segment the 12 nucleotide conserved influenza 3' terminal sequence has been modified by replacement of one to three nucleotides occurring in said
20 sequence at positions 3, 5 and 8 relative to the 3' end by other nucleotides provided that the nucleotides introduced in positions 3 and 8 are forming a base pair (i.e., if the nucleotide position 3 is G, than that in position 8 is C; if the nucleotide in position 3 is C, than that in position 8 is G; etc.).

25

The 3' conserved regions of the wild-type influenza virus have the following sequences:

Influenza A: (5')-CCUGCUUUUGCU-3'

Influenza B: (5')-NN(C/U)GCUUCUGCU-3'

30 Influenza C: (5')-CCUGCUUCUGCU-3'.

Moreover, the 13 nucleotide conserved influenza 5'-terminal sequence may be modified by replacement of one or two nucleotides occurring in said sequence as positions 3 and 8 by other nucleotides, again provided that the introduced nucleotides are forming a base pair. The 5' conserved regions of the wild-type influenza virus have the following sequences:

Influenza A: 5'-AGUAGAAACAAGG

Influenza B: 5'-AGUAG(A/U)AACA(A/G)NN

Influenza C: 5'-AGCAGUAGCAAG(G/A):

- 10 Preferred influenza viruses of the invention are those wherein in the 3' conserved region the replacements G3A and C8U have been performed, more preferred are those where also the replacement U5C has been performed (the above mutations are annotated relative to the 3' end; such counting from the 3' end is also indicated by a line on top of the digit, e.g.,
- 15 G $\overline{3}$ A). Another preferred influenza virus mutant comprises the 3'-terminal nucleotide sequence G3C, U5C and C8G (relative to the 3' end) resulting in the following 3' terminal nucleotide sequence (5')-CCUGGUUCUCCU-3'. Among the influenza viruses defined hereinbefore those having a 3'-terminal nucleotide sequence of (5')-CCUGUUUCUACU-3' are most
- 20 preferred. In case of an influenza A virus the segment may further have the modifications U3A and A8U in its 5' terminal sequence, in case of influenza C it may have the modifications C3U and G8A in its 5' terminal sequence. The most preferred influenza viruses of the present invention comprise the following general structures:

25 Influenza A (mutant pHL1104):

5'-AGUAGAAACAAGGNNNU₅₋₆..(880-2300 ntds)..N'N'N'CCUG \overline{U} UU \overline{C} U \overline{A} CU-3'

Influenza A (mutant pHL1920):

5'-AG \overline{A} AGAA \overline{U} CAAGGNNNU₅₋₆..(880-2300 ntds)..N'N'N'CCUG \overline{U} UU \overline{C} U \overline{A} CU-3'

Influenza A (mutant pHL1948):

- 30 5'-AGUAGAAACAAGGNNNU₅₋₆..(880-2300 ntds)..N'N'N'CCUG \overline{G} UU \overline{C} U \overline{C} CU-3'

Influenza B:

5'-AGUAG(A/U)AACA(A/G)NNNNNU₅₋₆..(880-2300 ntds)..N'N'N'N'(C/U)GUUUCUACU-3'

Influenza C:

5'-AGUAGUAACAAG(G/A)GU₅₋₆..(880-2300 ntds)..CCCCUGUUCUACU-3'

- 5 In the above structures the variables are defined as follows:
 - (1) Underlined and enlarged letters show the required mutations relative to the wild-type sequence for preparing a promoter mutant with enhanced properties;
 - (2) enlarged A in position 10 in the 5'-part of the sequence: unpaired A
 - 10 residue, bulge-forming;
 - (3) (A/G) in one position: different isolates or single segments with variable sequence at the respective position, which are functionally interchangeable;
 - (4) N and N': positions undefined, but base-paired relative to each other
 - 15 because of complementarity between the 5' and 3' termini, different among the 8 segments, but constant for each segment throughout all viral isolates;
 - (5) (880-2300 ntds): the lengths of the viral RNA segments, in case of segments with foreign genes increased up to 4,000 nucleotides.

20

According to embodiments (1) to (3) the invention provides

- a stable recombinant influenza virus containing (up to) seven regular vRNA segments plus one (or more) additional bicistronic segment(s) coding for a foreign gene in covalent conjunction with one of the influenza
- 25 genes, in tandem arrangement, and
- a method for the construction of stable recombinant influenza viruses through tandem arrangement of bicistronic vRNA segments, which is also applicable as a method for attenuation and for prevention of reassortment between co-infecting influenza viruses.

30

Expression of both gene products in these constructions is made possible by way of an upstream splice donor and a downstream splice acceptor

signal surrounding the proximal coding region of such a quality that splicing does occur in part of the mRNA molecules only, i.e., both mRNAs spliced and unspliced are present in the infected cell. For compensation with regard to the vRNA length the bicistronic segment is connected to a
5 promoter variant of enhanced replication and transcription rates as defined herein before.

The splice donor and the splice acceptor signals are selected from authentic sequences as present in influenza segments 7 and 8 or other
10 partially effective splice reaction substrates, preferably those of influenza virus WSN segment 7, i.e., 5'-AG[↓]GTACGTTC-3' (donor) and 5'-GCTGAAAAATGATCTTCTTGAAAATTGCAG[↓]GC-3' (acceptor).

In a particular application of embodiments (1) to (3) the tandem
15 bicistronic mRNA codes for one of the viral genes, such as hemagglutinin, in conjunction with all or part of the viral neuraminidase coding sequence, in antisense orientation, while the authentic neuraminidase vRNA segment is missing in these recombinant viruses. In another variation of these constructs an anti-neuraminidase ribozyme sequence is also provided
20 together with the (partial) neuraminidase antisense sequence, in the proximal position of these bicistronic recombinant segments. Recombinant viruses of this character are propagated in culture media with addition of exogenous neuraminidase.

25 The absence of a functional neuraminidase gene serves as a strong attenuation mechanism resulting in single-step infections of such recombinant viruses only. While a functional neuraminidase gene could be provided through another (wildtype) influenza virus superinfecting the same cell, expression of that gene is very much reduced through
30 antisense RNA interaction and/or destruction of the corresponding vRNA through ribozyme cleavage, designed to interfere with production of

infectious progeny even from co-infected cells; as a barrier against reassortment in double infected cells.

Recombinant viral RNAs coding simultaneously for two genes in tandem in
5 a construct in which one of the viral genes is in covalent junction with a
foreign coding sequence, are constructed via E. coli plasmid vector DNAs
designed for an *in vivo* transcription of minus-strand vRNAs by cellular
RNA polymerase I. In these constructs the gene in plus-strand proximal
(upstream) position is surrounded by splice signals of limited activity such
10 that both mRNAs, spliced and unspliced are present in the infected cell.
Either the foreign gene or the viral gene may be in that upstream position.
In the majority of applications the higher rates of expression will be
reserved for the foreign coding sequence, while the lower expression rate
of the viral gene is adapted to be approximately in balance with
15 expression of the other viral genes encoded by the regular viral segments.

To achieve such a balanced rate of expression, the splice signals and the
promoter have to be chosen properly (Flick and Hobom, Interaction of
influenza virus, polymerase with viral RNA in the 'corkscrew' conformation,
20 J. Gen. Virol. 80, 2565-2572 (1999)). At an increased overall transcription
rate, the resulting mRNAs shall be spliced inefficiently if the viral gene is in
the distal (downstream) position. Vice-versa, if the foreign gene is in the
distal position, splicing to obtain the foreign mRNA shall be achieved
efficiently. Both designs serve to reach an over-expression of the foreign
25 gene relative to the viral gene, of which the expression shall be in balance
with the expression of the other viral genes. Further, the promoter variant
attached to the bicistronic segment has the function to compensate for the
increased gene length by way of an increased replication rate.

30 The influenza vRNA segments preferably used for construction of
bicistronic segments include the neuraminidase (No. 6), hemagglutinin
(No. 4) and NS segment (No. 8). In the NS segment the foreign gene may

also substitute for the NS1 gene leaving the viral NS2 gene in its place. These recombinant viruses can, as an example, be made by the following procedure: A recombinant virus population can be selected by repeated ribozyme-mediated cleavage of helper-virus segments carrying ribozyme
5 cleavage sites that flank the same viral gene in the monocistronic segment as is present in the bicistronic construct (PCT/EP00/01903). By serial viral passaging and relying on the output of reporter genes in equivalently constructed bicistronic segments, a balanced mode of expression can be achieved in choosing the right set of elements: promoter, splice signals
10 plus a limited variation in segment length. The construct that gives rise to the balanced, stable expression is then used as a basis for a multiple cDNA transfection procedure in a helper-virus free design according to Neumann et al., Proc. Natl. Acad. Sci. USA, Vol 96, 9345-9350 (August 1999). The resulting recombinant influenza virus, obtained via single plaques in pure
15 helper-free state is subjected to another series of propagation steps to finally evaluate its properties.

In a particular application this design is used for a controlled mode of viral attenuation. Attenuation of influenza viruses so far has been achieved in
20 cold-sensitive mutants (Edwards et al., J. Infect. Dis. 169, 68-76(1994)), by deletion of the NS1 gene (partial attenuation, Egorov et al., J. Virol. 72, 6437-6441 (August 1998) and Palese et al., Proc. Natl. Acad. Sci USA, 4309-4314 (April 2000)), or through deletion of the neuraminidase gene (full attenuation, Kawaoka et al., J. Virol. 74, 5206-5212 (June 2000)).
25 The latter approach is adapted here using a novel technique for the attenuation, which for the first time is also able to interfere with (chance) superinfection by wild-type viruses.

In this embodiment of the invention a bicistronic cDNA construct is
30 achieved, which instead of a foreign gene is coding either for part of or for the entire viral neuraminidase gene in antisense orientation, with or without being surrounded both by splice donor and acceptor elements. In

another version of that design a 2 x 50 nucleotide antisense segment complementary to the 5'-terminal neuraminidase sequence has been cloned in flanking positions relative to a ribozyme construct according to the hammerhead design and oriented against a common GUC triplett
5 within the neuraminidase sequence. In a preferred design this antisense expression construct has been attached to the hemagglutinin vRNA segment, while another gene or reporter gene is encoded in a second bicistronic vRNA, in conjunction with NS2.

10 Propagation of recombinant viruses deleted for the neuraminidase (NA) gene requires an addition of external neuraminidase to the medium. In the absence of neuraminidase, infection by the NA deletion viruses is abortive: no infectious progeny is produced. Upon co-infection (3:3) of recombinant viruses together with wildtype viruses no progeny virus or plaque is
15 observed, which is attributed to antisense-blocked expression or (partial) destruction of the neuraminidase segment originating from the wild-type virus. Therefore, the recombinant viruses described are not only attenuated in single infections, but simultaneously interfere with wildtype virus superinfection, and therefore, no re-assortment between the two
20 viruses will occur.

The pharmaceutical composition according to embodiment (4) above and the medicament of embodiment (5) above contain the recombinant influenza virus in a pharmaceutically effective amount. Besides said
25 recombinant influenza virus, the pharmaceutical composition and the medicament may contain further pharmaceutically acceptable carrier substances well-known to a person skilled in the art, such as binders, desintegrants, diluents, buffers, preservatives, etc. The pharmaceutical composition and medicaments are solid or liquid preparations and are
30 suitable to be administered orally, intravenously or subcutaneously.

The medicament according to embodiment (5) above is preferably suitable as a medicament against influenza and/or against other infections. The recombinant influenza virus may be present in form of inactivated preparations or may be present in form of live recombinant viruses, preferably as attenuated viruses.

Live recombinant viral vaccines, live but attenuated recombinant viral vaccines or inactivated recombinant viral vaccine can be formulated. Inactivated vaccines are "dead" in the sense that their infectivity has been destroyed. Ideally, the infectivity is destroyed without affecting its immunogenicity. To prepare inactivated vaccines, the recombinant virus may be grown in cell cultures or in embryonated chicken eggs, purified, and inactivated by formaldehyde or β -propiolactone. The resulting vaccine is usually administered intramuscularly.

Inactivated viruses may be formulated with suitable adjuvants to enhance the immunological response. Such adjuvants include, but are not limited to, mineral gels, e.g., aluminum hydroxide, surface-active substances such as pluronic polyols, lysolecithin, peptides, oil emulsions, and potentially useful human adjuvants such as BCG.

Many methods may be used to introduce the vaccine formulations above, for example the oral, intradermal, intramuscular, intraperitoneal, subcutaneous, or intranasal routes. Where a live recombinant virus vaccine is used, it is preferred to introduce the formulation via the natural route of infection for influenza virus.

The medicament according to embodiment (5) above is preferably suitable for prophylactic or therapeutic vaccination, or both, against influenza and other infections. For example, recombinant viruses can be made for use in vaccines against HIV, hepatitis B virus, hepatitis C virus, herpes viruses, papilloma viruses, to name but a few. In one embodiment the

recombinant virus contains the genes for surface proteins of the viruses, in another the genes for non-structural or regulatory genes. The recombinant viruses may be present in form of inactivated preparations or may be present in form of live recombinant viruses, or as live, but
5 attenuated viruses. In an attenuated virus the recombinant virus would go through a single or at most very few propagation cycle(s) and induce a sufficient level of immune response, but would not cause disease. Such viruses lack one of the essential influenza genes or contain mutations to introduce temperature sensitivity.

10

The agents of embodiments (6)-(8) of the invention are applicable in *ex vivo* and *in vivo* application schemes. The RNA molecule to be expressed by means of the agent of the embodiment (8) is of an antisense sequence or double strand sequence (in ambisense bidirectional transcription)
15 relative to a target cellular mRNA molecule. In embodiment (8) the agent is preferably suitable for sequence-specific gene silencing, preferably by antisense RNA or RNA interference mechanisms.

20

The method for the production of proteins or glycoproteins is preferably performed in cell culture cells or in fertilized chicken cells in accordance with standard techniques within the general knowledge of a person skilled in the art. The proteins or glycoproteins to be expressed are those incorporated into the ambisense construct as defined hereinbefore.

25

The methods according to embodiments (9) to (12), (14) and (15) of the invention include the administration of an effective amount to the mammal or the administration of a sufficient infective dose of the recombinant virus to the cell system that is used for *ex vivo* therapy or for *in vitro* investigations, whereby the amount and dose will be determined by a
30 person skilled in the respective arts or knowledgeable of the desired treatments.

The agent of embodiment (14) of the invention is preferably utilized to infect, transfect or transduce patient-derived immune cells. The agent is suitable for treatment of cancer or chronic viral infections. For this purpose, patient derived immune cells, preferably dendritic cells, are ex vivo infected with recombinant influenza viruses expressing, e.g., tumor antigens or viral antigens. The transduced cells are then reintroduced into the patient.

The preferred method for Immunotherapy of embodiment (14) of the invention is an autologous immunotherapy, wherein the cells which are ex vivo infected are patient-derived and the transduced cells are reintroduced into the patient. The diseases to be treated by this method include cancer and chronic viral infections. For details regarding such treatment see discussion of embodiment (13) above.

The method for inducing antibodies according to embodiment (15) of the invention is suitable for inducing antibodies to foreign proteins including glycoproteins, following the administration of protein or glycoprotein antigens as part of a recombinant influenza virus in an authentic conformation, whereby the virus is purified by gentle procedures based on hemagglutination, and the gene is expressed at high rates in the infected cells.

As influenza viruses have a wide host range, recombinant influenza viruses can be used to obtain strong immune responses in, and isolate antibodies from, a wide range of animals, including, but not limited to, fowl, pigs, horses, and mice. Further, influenza viruses adapted to the mouse can be used for the infection of mice by several routes including the intranasal route. This results in infection of the pharyngeal mucosal cells and results in an additional type of B cell response (e.g., as recognized in the ratio of IgG to IgA). Mice are of particular utility in the induction of immune responses in transgenic mice that have been

engineered to express human antibodies. As gentle procedures based on hemadsorption are used to purify influenza viruses, antibodies to antigens in native conformation can be isolated from the infected mammals.

The present invention further illustrated by the following, non-limiting

5 Example:

Example

Model tandem bicistronic expression constructs using reporter genes CAT and GFP.

- 10 Objective: Measurements of relative expression rates for CAT in proximal and distal position, with live observation of GFP fluorescence in alternate position during propagation of recombinant influenza viruses.

a) Construction of bicistronic expression plasmid DNAs:

- Starting out with vector plasmid pHH10 (Hoffmann, Ph.D. Thesis, Univ. Giessen (1997)), i.e. an ampicillin resistant plasmid including in between a human rDNA promoter segment and a murine rDNA terminator segment precisely inserted cDNA sequence elements representing the 5' and 3' vRNA sequence of influenza rRNA segment 5, and finally a central multiple cloning site sequence as obtained from plasmid PBSK, both reporter genes
- 20 have been inserted in a stepwise manner. After that, to the proximal reading frame, i.e. CAT in pHL3196, and GFP in pHL3224, has been added an upstream splice donor sequence element and a downstream splice acceptor element, both inserted as double-strand oligonucleotides, in between particular restriction cleavage sites available in the respective
- 25 positions. The signal sequences used in that pair of plasmids indicated above have been derived from influenza vRNA segment 7, which is known for its partial splice reactions yielding both gene products, M1 and M2, simultaneously. By insertion of a non-transcribed DNA fragment (representing an internal segment of the influenza PB1 coding region) in a
- 30 distal position relative to both reading frames, pHL3196 has been converted into pHL3235, and pHL3224 into pHL3236. For the resulting plasmid constructs see Figs 5-8 and SEQ ID NOs: 21-24.

b) Transfection of plasmid DNAs and isolation of recombinant influenza viruses:

Semi-confluent 293-T cells, a human renal carcinoma cell line carrying an artificially integrated tumor virus SV40 T-antigen gene, were DNA-transfected using lipofectamine: 5-10 µg of DNA mixed with 30µl Lipofectamine® (GIBCO/BRL) were added to 370 µl of DMEM medium and were incubated with 5×10^6 to 10^7 cells, washed and maintained serum-free for 5 to 8 hours, before serum was added for another 12 to 15 hours. Finally influenza helper virus FPV_{Bratislava} was used for infection of the DNA-transfected cells. The supernatant containing a mixture of helper viruses and recombinant viruses was collected for further propagation after 8 to 12 hours of infection, while the sedimented cells were used for preparation of a cell lysate, fractions of which were inserted in the CAT assay procedure.

Viral propagation was achieved by infection of MDCK cells (Madin-Darby canine kidney cell line) again in semi-confluent state (5×10^6 to 10^7 cells per plate), generally using 1ml of the previous supernatant for infection. Serial propagations were done in the same way, with preparation of cell lysates for CAT assays at the end of each step. Infected cells were also used for observation of GFP fluorescence.

c) CAT assay:

Bacterial chloramphenicol-acetyltransferase (CAT) is accumulated in eukaryotic cells without degradation and can be used for representative gene expression measurements. The substrate used here is fluorescent boron-dipyrromethane-chloramphenicol difluoride (FLASH CAT-KIT®; Stratagene). 50 µl of cell lysate or reduced/diluted samples thereof were used for incubation with 7.5 µl of fluorescent substrate and 10 µl acetyl-CoA (4mM) co-substrate in 19 mM Tris/HCl, pH: 7.5 at 37°C for 3 hours. For extraction of reaction products 1ml of ethylacetate ins added, the

mixture is vortexed, and separated by centrifugation. After solvent evaporation and dissolution again in 20ml ethylacetate, the reaction products are separated on a silica thin-layer chromatography plate using chloroform/methanol 87:13% (vol.) and the results are documented by photography under UV light.

d) Results

CAT in proximal or in distal position of this pair of recombinant plasmids is expressed about equally (Fig.4), and the same is true for GFP (not shown). The expression rates are increasing during the initial steps of viral propagation and stay about constant afterwards during further steps of recombinant viral passages, different from expression rates in ambisense bicistronic constructs (pHL2899 and pHL2960) (Fig. 4B). Co-transfection of booster plasmids in the initial 293-T cells increase the yields of recombinant viruses within the progeny population, which are maintained during consecutive steps of propagation. Addition of 1000 nucleotides of untranslated vRNA sequence will not reduce the expression rates substantially (pHL3235 versus pHL3196, and pHL3236 versus pHL3224).

SEQUENCE LISTING

<110> ARTEMIS PHARMACEUTICALS GmbH

5 <120> Recombinant Influenza Viruses with Bicistronic vRNAs Coding for
Two Genes in Tandem Arrangement

<130> 011622wo/JH/ml

10 <140>
<141>

<160> 24

15 <170> PatentIn Ver. 2.1

<210> 1
<211> 12
<212> RNA

20 <213> Influenza A virus

<400> 1
ccugcuuuug cu 12

25

<210> 2
<211> 12
<212> RNA
<213> Influenza B virus

30

<400> 2
nnygcucucug cu 12

35 <210> 3
<211> 12
<212> RNA
<213> Influenza C virus

40 <400> 3
ccugcuucug cu 12

45 <210> 4
<211> 12
<212> RNA
<213> Artificial Sequence

<220>
50 <223> Description of Artificial Sequence: Modified
influenza A 3'-sequence (pHL1104 and pHL1920)

<400> 4

ccuguuucua cu 12

5 <210> 5
<211> 12
<212> RNA
<213> Artificial Sequence

10 <220>
<223> Description of Artificial Sequence: Modified
influenza A 3'-sequence (pHL1948)

15 <400> 5
ccugguucuc cu 12

20 <210> 6
<211> 13
<212> RNA
<213> Influenza A virus

<400> 6
aguagaaaca agg 13

25 <210> 7
<211> 13
<212> RNA
<213> Influenza B virus

30 <400> 7
aguagwaaca rnn 13

35 <210> 8
<211> 13
<212> RNA
<213> Influenza C virus

40 <400> 8
agcaguagca agr 13

45 <210> 9
<211> 13
<212> RNA
<213> Artificial Sequence

50 <220>
<223> Description of Artificial Sequence: Modified
influenza A 5'-sequence (pHL1920)

<400> 9

agaagaauca agg 13

5 <210> 10
<211> 21
<212> RNA
<213> Influenza A virus

10 <400> 10
aguagaaaca aggnnnuuuu u 21

15 <210> 11
<211> 21
<212> RNA
<213> Artificial Sequence

20 <220>
<223> Description of Artificial Sequence: Modified
influenza A 5'-sequence (pHL1920)

<400> 11
agaagaauca aggnnnuuuu u 21

25 <210> 12
<211> 21
<212> RNA
<213> Influenza B virus

30 <400> 12
aguagwaaca rnnnnuuuu u 21

35 <210> 13
<211> 19
<212> RNA
<213> Artificial Sequence

40 <220>
<223> Description of Artificial Sequence: Modified
influenza C 5'-sequence

45 <400> 13
aguaguaaca agrguuuuu 19

50 <210> 14
<211> 15
<212> RNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Modified
influenza A 3'-sequence (pHL1104 and pHL1920)

<400> 14
5 nnnccuguuu cuacu 15

<210> 15
<211> 15
10 <212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Modified
15 influenza A 3'-sequence (pHL1948)

<400> 15
nnnccugguu cuccu 15

20 <210> 16
<211> 15
<212> RNA
<213> Artificial Sequence

25 <220>
<223> Description of Artificial Sequence: Modified
influenza B 3' sequence

30 <400> 16
nnnnnyguuu cuacu 15

35 <210> 17
<211> 14
<212> RNA
<213> Artificial Sequence

40 <220>
<223> Description of Artificial Sequence: Modified
influenza C 3'-sequence

<400> 17
45 ccccuguuuc uacu 14

<210> 18
<211> 10
<212> DNA
50 <213> Influenza A virus

<400> 18
aggtacgttc 10

<210> 19
 <211> 32
 5 <212> DNA
 <213> Influenza A virus

<400> 19
 10 gctgaaaaat gatcttcttg aaaattgcag gc 32

<210> 20
 <211> 3888
 <212> DNA
 15 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: pHL1920

20 <400> 20
 cccaaaaaaaa aaaaaaaaaa aaaaaaaaaag agtccagagt ggccccgccg ttccgcgcgcg 60
 ggggggggggg ggggggggga cacttttcgga catctggtcg acctccagca tcgggggaaa 120
 aaaaaaaaaac aaagttttcgc ccggagtact ggtcgacctc cgaagttggg ggggagtaga 180
 aacagggttag ataatcactc actgagtgcac atccacatcg cgagcgcgcg taatacgact 240
 25 cactataggg cgaattgggt accgggcccc ccctcgaggt cgacggtatc gataagcttc 300
 gacgagattt tcaggagcta aggaagctaa aatggagaaa aaaatcactg gatataccac 360
 cgttgatata tcccaatggc atcgtaaaga acattttgag gcatttcagt cagttgctca 420
 atgtacctat aaccagaccg ttcagctgga tattacggcc tttttaaaga ccgtaaagaa 480
 aaataagcac aagttttatc cggcctttat tcacattctt gccgcctga tgaatgctca 540
 30 tccggaattc cgtatggcaa tgaaagacgg tgagctggtg atatgggata gtgttcaccc 600
 ttgttacacc gttttccatg agcaaaactga aacgttttca tcgtctgga gtgaatacca 660
 cgacgatttc cggcagtttc tacacatata ttcgcaagat gtggcgtgtt acggtgaaaa 720
 cctggcctat ttccctaaag ggtttattga gaatatgttt ttctctcag ccaatccctg 780
 ggtgagtttc accagttttg atttaaactg ggccaatatg gacaacttct tcgcccccg 840
 35 tttcaccatg ggcaaataat atacgcaagg cgacaaggtg ctgatgccgc tggcgattca 900
 ggttcatcat gccgtttgtg atggcttcca tgtcggcaga atgcttaatg aattacaaca 960
 gtactgcgat gagtggcagg gcggggcgta atttttttaa ggcagttatt ggtgccctta 1020
 aacgcctggt gctacgcctg aataagtgat aataagcggg tgaatggcag aaattcgctc 1080
 aagcttgata tcgaattcct gcagcccggg ggatccacta gttctagagc ggccgccacc 1140
 40 gcggtggagc tccagctttt gttcccttta gtgagggtta attgcgcgca ggcctagcta 1200
 ggtaaagaaa aatacccttg attcttctaa taaccggcg gccccaaatg ccgactcgga 1260
 gcgaaagata tacctcccc ggggcccggga ggtcgcgtca ccgaccacgc cgccggccca 1320
 ggcgacgcgc gacacggaca cctgtcccca aaaacgccac catcgcagcc acacacggag 1380
 cgcccggggc cctctggtca accccaggac acacgcggga gcagcgcgcg gccggggacg 1440
 45 ccctcccggc cgcccgtgcc acacgcaggg ggccggccc tgtctccaga gcgggagccg 1500
 gaagcatttt cgccgggccc ctccctacgac cgggacacac gagggaccga aggcgggcca 1560
 ggcgcgacct ctggggccgc acgcgcgctc agggagcgtc ctccgactcc gcacggggac 1620
 tcgcagaaa ggatcgtgac ctgcattaat gaatcagggg ataacgcagg aaagaacatg 1680
 tgagcaaaaag gccagcaaaa ggccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc 1740
 50 cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 1800
 aaccggacag gactataaag ataccaggcg tttcccctg gaagctccct cgtgcgctct 1860
 cctgttccga ccctgcgcgt tacgggatac ctgtccgcct ttctcccttc gggaagcgtg 1920
 gcgctttctc atagctcacg ctgtaggtat ctgagttcgg ttaggtcgt tcgctccaag 1980

31

	ctgggctgtg	tgcacgaacc	ccccgttcag	cccgcaccgt	gcgccttata	cggtaactat	2040
	cgtcttgagt	ccaacccggg	aagacacgac	ttatcgccac	tggcagcagc	cactggtaac	2100
	aggattagca	gagcgaggta	tgtaggcggg	gctacagagt	tcttgaagtg	gtggcctaac	2160
	tacggctaca	ctagaaggac	agtatttggg	atctgcgctc	tgctgaagcc	agttaccttc	2220
5	ggaaaaagag	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	cgggtggtttt	2280
	tttgtttgca	agcagcagat	tacgcgcaga	aaaaaaggat	ctcaagaaga	tcctttgatc	2340
	ttttctacgg	ggctctgacgc	tcagtggaaac	gaaaactcac	gttaagggat	tttgggtcatg	2400
	agattatcaa	aaaggatctt	cacctagatc	cttttaaatt	aaaaatgaag	ttttaaatca	2460
	atctaaagta	tatatgagta	aacttgggtc	gacagttacc	aatgcttaat	cagtgaggca	2520
10	cctatctcag	cgatctgtct	atttcgttca	tcctagttg	cctgactccc	cgtcgtgtag	2580
	ataactacga	tacgggaggg	cttaccatct	ggccccagtg	ctgcaatgat	accgcgagac	2640
	ccacgctcac	cggctccaga	tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc	2700
	agaagtgggc	ctgcaacttt	atccgcctcc	atccagtcta	ttaattgttg	ccgggaagct	2760
	agagtaagta	gttcgccagt	taatagtttg	cgcaacgttg	ttgccattgc	tacaggcatc	2820
15	gtgggtgtcac	gctcgtcggt	tggtatggct	tcattcagct	ccggttccca	acgatcaagg	2880
	cgagttacat	gatcccccat	gttggtgcaa	aaagcgggta	gctccttcgg	tcctccgatc	2940
	gttgctcagaa	gtaagttggc	cgcagtgtta	tcactcatgg	ttatggcagc	actgcataat	3000
	tctcttactg	tcattgccatc	cgtaagatgc	ttttctgtga	ctgggtgagta	ctcaaccaag	3060
	tcattctgag	aatagtgtat	gcggcgaccg	agttgctctt	gcccggcgtc	aacacgggat	3120
20	aataccgcgc	cacatagcag	aactttaaaa	gtgctcatca	ttggaaaacg	ttcttcgggg	3180
	cgaaaaactct	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	cactcgtgca	3240
	cccaactgat	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga	3300
	aggcaaaatg	ccgcaaaaaa	gggaataagg	gcgacacgga	aatggtgaat	actcatactc	3360
	ttcctttttc	aatattattg	aagcatttat	cagggttatt	gtctcatgag	cggatacata	3420
25	tttgaatgta	tttagaaaaa	taaaacaaaag	agtttgtaga	aacgcaaaaa	ggccatccgt	3480
	caggatggcc	ttctgcttaa	ttgatgcct	ggcagtttat	ggcgggcgtc	ctgcccgcca	3540
	ccctccgggg	cggtgcttcg	caacgttcaa	atccgctccc	ggcggatttg	tcctcccgca	3600
	gagagcggtc	accgacaaac	aacagataaa	acgaaaaggc	cagtctttcg	actgagcctt	3660
	tcgttttatt	tgatgcctgg	cagttcccta	ctctcgcatg	gggagacccc	acactaccat	3720
30	cggcgctacg	gcgtttcact	tctgagttcg	gcattggggtc	aggtgggacc	accgcgctac	3780
	tgccgccagg	caaattctgt	tttatcagac	cgcttctgcg	ttctgattta	atctgtatca	3840
	ggctgaaaaat	cttctctcat	ccgccaaaac	agaagctagc	ggccgatc		3888
35	<210> 21						
	<211> 4500						
	<212> DNA						
	<213> Artificial Sequence						
40	<220>						
	<223> Description of Artificial Sequence: pHL3196						
	<400> 21						
	agtagaaaca	gggtagataa	tcactcactg	agtgacatcc	acatcgcgag	cgcgagggta	60
45	cgttctcgag	cgcgcgtaat	acgactcact	atagggcgaa	ttgggtacgt	tcctcatatg	120
	agaaaaaaat	cactggatat	accaccgttg	atatatccca	atggcatcgt	aaagaacatt	180
	ttgaggcatt	tcagtcagtt	gctcaatgta	cctataacca	gaccgttcag	ctggatatta	240
	cggccttttt	aaagaccgta	aagaaaaata	agcacaagtt	ttatccggcc	tttattcaca	300
	ttcttgcccc	cctgatgaat	gctcatccgg	aattccgtat	ggcaatgaaa	gacgggtgagc	360
50	tggtgatatg	ggatagtgtt	cacccttggt	acaccgtttt	ccatgagcaa	actgaaacgt	420
	tttcatcgct	ctggagtga	taccacgacg	atttccggca	gtttctacac	atatattcgc	480
	aagatgtggc	gtgttacggg	gaaaacctgg	cctatttccc	taaaggggtt	attgagaata	540
	tgtttttcgt	ctcagccaat	ccctgggtga	gtttcaccag	ttttgattta	aacgtggcca	600

	atatggacaa	cttcttcgcc	cccgttttca	ccatgggcaa	atattatacg	caaggcgaca	660
	aggtgctgat	gccgctggcg	attcaggttc	atcatgccgt	ctgtgatggc	ttccatgtcg	720
	gcagaatgct	taatgaatta	caacagtact	gcgatgagtg	gcagggcggg	gcgcgttaac	780
	gagatcagct	gaaaaatgat	cttcttgaaa	atgtgcaggc	cgtacgtgta	ccgggcccc	840
5	cctcgactcg	cgaaggagtc	caccatgagt	aaaggagaag	aacttttcac	tggagtgtgc	900
	ccaattcttg	ttgaattaga	tggatgatgt	aatgggcaca	aattttctgt	cagtggagag	960
	ggtgaagggtg	atgcaacata	cggaaaactt	acccttaa	ttatttgcac	tactggaaaa	1020
	ctacctgttc	catggccaac	acttgtcact	actttcactt	atggtgttca	atgcttttca	1080
	agatacccag	atcatatgaa	acagcatgac	tttttcaaga	gtgccaatgcc	cgaagggtat	1140
10	gtacaggaaa	gaactatatt	tttcaaaagat	gacgggaact	acaagacacg	tgctgaagtc	1200
	aagtttgaag	gtgataccct	tgtaataaga	atcgagttaa	aaggatttga	ttttaaaaga	1260
	gatggaaaca	ttcttggaca	caaattggaa	tacaactata	actcacacaa	tgatacatc	1320
	atggctgaca	agcagaagaa	cggaaatcaag	gccaacttca	agaccgcga	caacatcgag	1380
	gacggcgggc	tgcatgtggc	cgaccactac	cagcagaaca	cccccaattg	cgatggccct	1440
15	gtcctttttc	cagacaacca	ttacctgtcc	acacaatctg	ccctttcgaa	agatcccaac	1500
	gaaaagagag	accacatggt	ccttcttgag	tttgtaacag	ctgctgggat	tacacatggc	1560
	atggatgaac	tatacaagg	atcccatcac	catcaccatc	actaagctcc	atggtctaga	1620
	tatcgatagg	cctagctagg	taaagaaaaa	tacccttggt	tctactaata	acccggcggc	1680
	ccaaaatgcc	gactcggagc	gaaagatata	cctcccccg	ggccgggagg	tcgcgtcacc	1740
20	gaccacgccg	ccggcccagg	cgacgcgcga	cacggacacc	tgccccaaa	aacgccacca	1800
	tcgcagccac	acacggagcg	cccggggccc	tctgggtcaac	cccaggacac	acgcgggagc	1860
	agcgccgggc	cggggacgcc	ctcccggccg	cccgtgccac	acgcaggggg	ccggcccgtg	1920
	tctccagagc	gggagccgga	agcattttcg	gcccggccct	cctacgaccg	ggacacacga	1980
	gggaccgaag	gccggccagg	cgcgacctct	cgggcccgcac	gcgcgctcag	ggagcgctct	2040
25	ccgactccgc	acggggactc	gccagaaagg	atcgtgacct	gcattaatga	atcaggggat	2100
	aacgcaggaa	agaacatgtg	agcaaaaggc	cagcaaaagg	ccaggaaccg	taaaaaggcc	2160
	gcgttgctgg	cgttttttcca	taggctccgc	ccccctgacg	agcatcaca	aaatcgacgc	2220
	tcaagtcaga	ggtggcgaaa	cccagacagga	ctataaagat	accaggcggt	ttcccctgga	2280
	agctccctcg	tgcgctctcc	tggtccgacc	ctgcccgtta	ccggatacct	gtccgccttt	2340
30	ctcccttcgg	gaagcgtggc	gctttctcat	agctcacgct	gtaggtatct	cagttcgggtg	2400
	taggtcgttc	gctccaagct	gggctgtgtg	cacgaacccc	ccgttcagcc	cgaccgctgc	2460
	gccttatccg	gtaactatcg	tcttgagtcc	aaaccggtaa	gacacgactt	atcgccactg	2520
	gcagcagcca	ctggtaacag	gattagcaga	gcgaggatag	taggcgggtg	tacagagttc	2580
	ttgaagtgg	ggcctaacta	cggctacact	agaaggacag	tatttggtat	ctgcgctctg	2640
35	ctgaagccag	ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc	2700
	gctggtagcg	gtggtttttt	tggttgcaag	cagcagatta	cgcgcagaaa	aaaaggatct	2760
	caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	agtggaaacg	aaactcacgt	2820
	taagggattt	tggtcatgag	attatcaaaa	aggatcttca	cctagatcct	tttaaattaa	2880
	aaatgaagtt	ttaaatcaat	ctaaagtata	tatgagtaaa	cttggtctga	cagttaccaa	2940
40	tgcttaatac	gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcatc	catagttgcc	3000
	tgactccccg	tcgtgtagat	aactacgata	cgggagggct	taccatctgg	ccccagtgct	3060
	gcaatgatac	cgcgagaccc	acgctcaccg	gctccagatt	tatcagcaat	aaaccagcca	3120
	gccggaagg	ccgagcgcag	aagtgtcctt	gcaactttat	ccgcctccat	ccagtctatt	3180
	aattgttgcc	gggaagctag	agtaagtagt	tcgccagtta	atagtttgcg	caacgttggt	3240
45	gccattgcta	caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	attcagctcc	3300
	gggtcccaac	gatcaaggcg	agttacatga	ttcccctatg	tggtgcaaaa	agcgggttagc	3360
	tccttcgggtc	ctccgatcgt	tgtcagaagt	aagttggccg	cagtgttatc	actcatgggt	3420
	atggcagcac	tgcataattc	tcttactgtc	atgccatccg	taagatgctt	ttctgtgact	3480
	ggtgagtagt	caaccaagtc	attctgagaa	tagtgtatgc	ggcgaccgag	ttgctcttgc	3540
50	ccggcgctca	cacgggataa	taccgcgcga	catagcagaa	ctttaaaagt	gctcatcatt	3600
	ggaaaacggt	cttcggggcg	aaaactctca	aggatcttac	cgctgttgag	atccagttcg	3660
	atgtaaccca	ctcgtgcacc	caactgatct	tcagcatctt	ttactttcac	cagcgtttct	3720
	gggtgagcaa	aaacaggaag	gcaaaatgcc	gcaaaaaagg	gaataagggc	gacacggaaa	3780

33

5 tgggtgaatac tcatactcctt ccttttttcaa tattattgaa gcatttatca gggttattgt 3840
 ctcatgagcg gatacatatt tgaatgtatt tagaaaaata aacaaaagag ttgttagaaa 3900
 cgcaaaaagg ccatccgtca ggatggcctt ctgcttaatt tgatgcctgg cagtttatgg 3960
 cgggcgctcct gcccgccacc ctccgggccc ttgcttcgca acgttcaa ccgtcccg 4020
 cggatttgtc ctactcagga gagcggtcac cgacaaacaa cagataaaac gaaaggccca 4080
 gtctttcgac tgagcctttc gttttatttg atgcctggca gttccctact ctgcgatggg 4140
 gagacccac actaccatcg gcgctacggc gtttcacttc tgagttcggc atggggtcag 4200
 gtgggaccac cgcgctactg ccgccaggca aattctgttt tatcagaccg cttctgcgtt 4260
 ctgatttaat ctgtatcagg ctgaaaatct tctctcatcc gccaaaacag aagctagcgg 4320
 10 ccgatcccca aaaaaaaaaa aaaaaaaaaa aaaaagagtc cagagtggcc ccgccgttcc 4380
 gcgccggggg gggggggggg ggggggacact ttcggacatc tggtcgacct ccagcatcgg 4440
 gggaaaaaaa aaaaacaaag tttcgcccg agtactggtc gacctccgaa gttggggggg 4500

15 <210> 22
 <211> 4721
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> Description of Artificial Sequence: pHL3224

 <400> 22
 25 atctagacca tggagcttag tgatggtgat ggtgatggga tcccttgat agttcatcca 60
 tgccatgtgt aatcccagca gctgttacaa actcaagaag gaccatgtgg tctctctttt 120
 cgttgggagc tttcgaaagg gcagattgtg tggacaggta atggttgctc ggtaaaagga 180
 cagggccatc gccaatggg gtgttctgct ggtagtggtc ggccagctgc acgccgccgt 240
 cctcgatgtt gtggcgggtc ttgaagtgg ccttgattcc gttcttctgc ttgtcagcca 300
 tgatgtatac attgtgtgag ttatagtgt attccaattt gtgtccaaga atgtttccat 360
 30 cttctttaaa atcaatacct ttttaactcg ttctattaac aagggtatca ccttcaaaact 420
 tgacttcagc acgtgtcttg tagttcccg catctttgaa aaatatagtt ctttctgtga 480
 cataaccttc gggcatggca ctcttgaaaa agtcatgctg tttcatatga tctgggtatc 540
 ttgaaaagca ttgaacacca taagtgaag tagtgacaag tgttggccat ggaacaggta 600
 gttttccagt agtgcaaata aatttaagg taagttttcc gtatgttgca tcaccttcac 660
 35 cctctccact gacagaaaat ttgtgcccat taacatcacc atctaattca acaagaattg 720
 ggacaactcc agtgaaaagt tcttctcctt tactcatggt ggactccttc gcgagtcgag 780
 ggggggccc gtacacgtac gcgctcgaga acgtaccttc gcgctcgaga tgtggatgtc 840
 actcagtgag tgattatcta ccctgtttct actcccccc aacttcggag gtcgaccagt 900
 actccggcgc aaactttgtt tttttttttt cccccgatgc tggaggtcga ccagatgtcc 960
 40 gaaagtgtcc cccccccccc ccccccccg cgcggaacgg cggggccact ctggactcct 1020
 tttttttttt tttttttttt ttttggggat cgcccgctag cttctgtttt ggcggatgag 1080
 agaagatttt cagcctgata cagattaaat cagaacgcag aagcggctctg ataaaacaga 1140
 atttgcctgg cggcagtagc gcggtgggtc cacctgaccc catgccgaac tcagaagtga 1200
 aacgccgtag cgccgatggg agtgtgggtc ctcccatg gagagtaggg aactgccagg 1260
 45 catcaataaa aacgaaaggc tcagtcgaaa gactgggcct ttcgttttat ctgttggttg 1320
 tcggtgaacg ctctcctgag taggacaaat ccgcccggag cggatttgaa cgttcggaag 1380
 caacggcccc gaggggtggc ggcaggacgc ccgccataaa ctgccaggca tcaaattaag 1440
 cagaaggcca tctgacgga tggccttttt gcgtttctac aaactctttt gtttattttt 1500
 ctaaatacat tcaaataatgt atccgctcat gagacaataa cctgataaaa tgcttcaata 1560
 50 atattgaaaa aggaagagta tgagtattca acatttccgt gtcgccctta ttcctttttt 1620
 tgcggcattt tgccttctctg tttttgctca ccagaaacg ctggtgaaag taaaagatgc 1680
 tgaagatcag ttgggtgcac gagtgggtta catcgaaactg gatctcaaca gcgtaagat 1740
 ccttgagagt tttcgccccg aagaacgttt tccaatgatg agcactttta aagttctgct 1800

	atgtggcgcg	gtattatccc	gtgttgacgc	cgggcaagag	caactcggtc	gccgcataca	1860
	ctattctcag	aatgacttgg	ttgagtactc	accagtcaca	gaaaagcatc	ttacggatgg	1920
	catgacagta	agagaattat	gcagtgcctg	cataaccatg	agtataaca	ctggcgccaa	1980
	cttacttctg	acaacgatcg	gaggaccgaa	ggagctaac	gcttttttgc	acaacatggg	2040
5	ggatcatgta	actcgccttg	atcgttggga	accggagctg	aatgaagcca	taccaaacga	2100
	cgagcgtgac	accacgatgc	ctgtagcaat	ggcaacaacg	ttgcgcaaac	tattaactgg	2160
	cgaactactt	actctagctt	cccggaaca	attaatagac	tggatggagg	cggataaagt	2220
	tgcaggacca	cttctgcgct	cggcccttcc	ggctggctgg	tttattgctg	ataaactctg	2280
	agccgggtgag	cgtgggtctc	gcggtatcat	tgcagcactg	gggacagatg	gtaagccctc	2340
10	ccgtatcgtg	gttatctaca	cgacggggag	tcaggcaact	atggatgaac	gaaatagaca	2400
	atatatactt	tagattgatt	taaaacttca	gcattggtaa	ctgtcagacc	aagtttactc	2460
	cctttttgat	aatctcatga	ccaaaatccc	tttaacgtgag	ttttcgttcc	actgagcgtc	2520
	agaccccgta	gaaaagatca	aaggatcttc	ttgagatcct	ttttttctgc	gcgtaactctg	2580
15	ctgcttgcaa	acaaaaaac	caccgctacc	agcggtggtt	tgtttgccgg	atcaagagct	2640
	accaactctt	tttccgaagg	taactggctt	cagcagagcg	cagataccaa	atactgtcct	2700
	tctagtgtag	ccgtagttag	gccaccactt	caagaactct	gtagcaccgc	ctacatacct	2760
	cgctctgcta	atcctgttac	cagtggctgc	tgccagtggc	gataagtcgt	gtcttaccgg	2820
	gttgactca	agacgatagt	taccggataa	ggcgcagcgg	tcgggctgaa	cgggggggtc	2880
20	gtgcacacag	cccagcttgg	agcgaacgac	ctacaccgaa	ctgagatacc	tacagcgtga	2940
	gctatgagaa	agcgcacgac	ttcccgaagg	gagaaaggcg	gacaggtatc	cggtaagcgg	3000
	cagggtcgga	acaggagagc	gcacgaggga	gcttccaggg	ggaaacgcct	ggtatcttta	3060
	tagtcctgtc	gggtttcgcc	acctctgact	tgagcgtcga	tttttgtgat	gctcgtcagg	3120
	ggggcgaggc	ctatggaaaa	acgccagcaa	cgcggccttt	ttacggttcc	tggccttttg	3180
25	ctggcctttt	gctcacatgt	tctttcctgc	gttatccctt	gattcattaa	tgcaggtcac	3240
	gaccccttct	ggcgagctcc	cgtgcggagt	cggagagcgc	tccctgagcg	cgcgtgcggc	3300
	ccgagaggtc	gcgcctggcc	ggccttcggt	ccctcgtgtg	tcccggtcgt	aggagggggc	3360
	ggccgaaaaa	gcttccggct	ccgctctggg	agacacgggc	cggccccctg	cgtgtggcac	3420
	ggggcgccgg	gagggcgctc	ccggcccgcc	gctgctcccg	cgtgtgtcct	gggttgacc	3480
30	agaggggccc	ggggcgctcc	tgtgtggctg	cgatggtggc	gtttttgggg	acaggtgtcc	3540
	gtgtcgcgcg	tcgcctgggc	cggcgcgctg	gtcggtgacg	cgacctcccg	gccccggggg	3600
	aggtatatct	ttcgctccga	gtcggcattt	tgggcccggc	ggttattagt	agaaacaagg	3660
	gtatttttct	ttacctagct	aggcctgcgc	gcaattaacc	ctcactaaag	ggaacaaaag	3720
	ctggagctcc	accgcggtgg	cggcgcctct	agaactagtg	gatcccccg	gctgcaggaa	3780
35	ttcgatatca	agcttcgacg	aattttctgcc	attcatccgc	ttattatcac	ttattcaggc	3840
	gtagcaccag	gcgtttaagg	gcaccaataa	ctgccttaaa	aaaattacgc	cccgccctgc	3900
	cactcatcgc	agtactgttg	taattcatta	agcattctgc	cgacatggaa	gccatcacaa	3960
	acggcatgat	gaacctgaat	cgccagcggc	atcagcacct	tgtcgccttg	cgtataatat	4020
	ttgcccatgg	tgaaaacggg	ggcgaagaag	ttgtccatat	tggccacggt	taaatcaaaa	4080
40	ctggtgaaac	tcacccaggg	attggctgag	acgaaaaaca	tattctcaat	aaacccttta	4140
	gggaaatagg	ccagggttttc	accgtaacac	gccacatctt	gcgaatatat	gtgtagaaac	4200
	tgccggaaat	cgtcgtggta	ttcactccag	agcgatgaaa	acgtttcagt	ttgtcatggg	4260
	aaaacgggtg	aacaagggtg	aacactatcc	catatcacca	gctcaccgtc	tttcattgcc	4320
	atacgaatt	cgggatgagc	attcatcagg	cgggcaagaa	tgtgaataaa	ggccggataa	4380
45	aacttgtgct	tatttttctt	tacggtcttt	aaaaaggccg	taatatccag	ctgaacggtc	4440
	tgggttatagg	tacattgagc	aactgactga	aatgcctcaa	aatgttcttt	acgatgccat	4500
	tgggatatat	caacgggtgg	atatccagtg	atttttttct	ccatttttagc	ttccttagct	4560
	cctgaaaatc	tcgtcgaagc	ttatcgatac	cgtcgacctc	gagggggggc	ccggtacggc	4620
	ctgcaaattt	tcaagaagat	cattttttcag	ctgatctcgt	t		4680
50							4721

<210> 23

<211> 5517

<212> DNA

<213> Artificial Sequence

<220>

5 <223> Description of Artificial Sequence: pHL3235

<400> 23

```

agtagaaaca gggtagataa tcaactcact agtgacatcc acatcgcgag cggaaggta 60
cgttctctag cgcgcgtaac acgactcact atagggcgaa ttgggtacgt tccatcatgg 120
10 agaaaaaaat cactggatat accaccgttg atatatccca atggcatcgt aaagaacatt 180
ttgaggcatt tcagtcagtt gctcaatgta cctataacca gaccgttcag ctggatatta 240
cggccttttt aaagaccgta aagaaaaata agcacaagtt ttatccggcc ttatttcaca 300
ttcttgcccg cctgatgaat gtcctccgg aattccgtat ggcaatgaaa gacggtgagc 360
tggtgatatg ggatagtgtt cacccttggt acaccgtttt ccatgagcaa actgaaacgt 420
15 tttcatcgct ctggagtga taccacgacg atttccggca gtttctacac atatattcgc 480
aagatgtggc gtgttacggt gaaaacctgg cctatttccc taaagggttt attgagaata 540
tggttttctg ctacagcaat ccctgggtga gtttcaccag ttttgattta aacgtggcca 600
atatggacaa cttcttcgcc ccctgtttca ccatgggcaa atattatacg caaggcgaca 660
aggtgctgat gccgctggcg attcagggtc atcatgccgt ctgtgatggc ttccatgtcg 720
20 gcagaatgct taatgaatta caacagtact gcgatgagtg gcagggcggg gcgcgttaac 780
gagatcagct gaaaaatgat cttcttgaaa atttcaggc cgtacgtgta cgggcccccc 840
cctcgactcg cgaaggagtc caccatgagt aaaggagaag aacttttcac tggagtgtgc 900
ccaattcttg ttgaattaga tggatggtt aatgggcaca aattttctgt cagtggagag 960
gggtgaagggtg atgcaacata cggaaaactt acccttaaat ttatttgcac tactggaaaa 1020
25 ctacctgttc catggccaac acttgtcact actttcactt atggtgttca atgcttttca 1080
agatacccag atcatatgaa acagcatgac tttttcaaga gtgcatgcc cgaagggtat 1140
gtacaggaaa gaactatatt tttcaaagat gacgggaact acaagacacg tgctgaagtc 1200
aagtttgaag gtgataccct tgttaataga atcgagttaa aaggattga ttttaaagaa 1260
gatggaaaca ttcttgga caaattggaa tacaactata actcacaca tgtatacatc 1320
30 atggctgaca agcagaagaa cggaatcaag gccaaacttca agaccgcga caacatcgag 1380
gacggcgggc tgcagctggc cgaccactac cagcagaaca cccaattgg cgatggccct 1440
gtccttttac cagacaacca ttacctgtcc acacaatctg ccctttcgaa agatcccaac 1500
gaaaagagag accacatggt cttcttgag tttgtaacag ctgctgggat tacacatggc 1560
atggatgaac tatacaaggg atcttcatga tctcagcaaa ctcttccttc ttaatccttc 1620
35 cagactcgaa gtcaattcgt gcacaaatcc gggccctaga caccatggcc tccaccatac 1680
tggaatttcc aactggctct ctgtatgagc tgctagggaa gaatttctcg aatagggtgc 1740
aacacttctg gtacatttgt tcatcctcaa ggattccctt ttgactcgta ttgagaatgg 1800
aacggtttct cttagggatc caagagtgtg tagttgccac agcatcatat tccatgcttt 1860
tggtctggacc atgggctggc attaccgcag cattgtttac agattcaatt tccttatgac 1920
40 tgacaaacgg gtatcatgga ttacaaagtc ttccctgata gtcttcatcc attagtccc 1980
atctcaggca aacttcggg atgtggagat tccgaatgt gtacagggtt ggcccgccat 2040
ctgaaaccaa cagtcctgcc tttgagcggg tctgtcccca cagcttcttt agctcgaatg 2100
acctcctcgt ttggatttgt gtgtctcccc tgtgacaccg gtatgtatat ctgtatcct 2160
tgatgaataa ttggagagcc atttgggctg ttgcccgtcc aagatcattg tttatcatgt 2220
45 tattctttat cactgttact ccaatgctca tatcagccga ttcattaatt cctgatactc 2280
caaagctggg caactccata ctaaaattgg ctacaaatcc atagcggtag aaaaagcttg 2340
tgaattcgaa tgttcctgtc ctatttatat aggacttttt ctgtctcata ttgatcccaa 2400
ctagcttgca ggttctgtag aatctatcca ctcccgcttg tattccctca tgatttggtg 2460
cattcacgat gagagcaaaa tcatcagagg actgaagtcc atcccaccag tatgtggttt 2520
50 tgggtgatct cttttgcca agattcagga ttgagactcc caacactgta ctacagatgt 2580
tgaacatacc catcatcatt cccgggctta atgaggctgt gccgtctatt atgagaggat 2640
cgataggcct agctaggtaa agaaaaatac ccttgtttct actaataacc cggcgcccca 2700
aaatgccgac tcggagcgaa agatatacct ccccgggggc cgggaggtcg cgtcaccgac 2760

```

	cacgccgccc	gcccaggcga	cgcgcgacac	ggacacctgt	ccccaaaaac	gccaccatcg	2820
	cagccacaca	cggagcgccc	ggggccctct	ggtcaacccc	aggacacacg	cgggagcagc	2880
	gccgggccc	ggacgcctc	ccggccgccc	gtgccacacg	cagggggccg	gcccgtgtct	2940
	ccagagcggg	agccggaagc	attttcggcc	ggccccctct	acgaccggga	cacacgaggg	3000
5	accgaaggcc	ggccaggcgc	gacctctcgg	gccgcacgcg	cgctcagggg	gcgctctccg	3060
	actccgcacg	gggactcgcc	agaaaggatc	gtgacctgca	ttaatgaatc	aggggataac	3120
	gcaggaaaga	acatgtgagc	aaaaggccag	caaaaggcca	ggaaccgtaa	aaaggccgcg	3180
	ttgctggcgt	ttttccatag	gctccgcccc	cctgacgagc	atcacaaaaa	tcgacgctca	3240
	agtacagagg	ggcgaaaccc	gacaggacta	taaagatacc	aggcgtttcc	ccctggaagc	3300
10	tccctcgtgc	gctctcctgt	tccgaccctg	ccgcttaccg	gatacctgtc	cgcttttctc	3360
	ccttcgggaa	gcgtggcgct	ttctcatagc	tcacgctgta	ggtatctcag	ttcgggtgtg	3420
	gtcgttcgct	ccaagctggg	ctgtgtgcac	gaaccccccg	ttcagcccga	ccgctgcgcc	3480
	ttatccggta	actatcgtct	tgagtccaac	ccggtaagac	acgacttatc	gccactggca	3540
	gcagccactg	gtaacaggat	tagcagagcg	aggatgtgag	gcgggtgctac	agagttcttg	3600
15	aagtgggtgg	ctaactacgg	ctacactaga	aggacagtat	ttggtatctg	cgctctgctg	3660
	aagccagtta	ccttcggaaa	aagagttggg	agctcttgat	ccggcaaaaa	aaccaccgct	3720
	ggtagcgggtg	gtttttttgt	ttgcaagcag	cagattacgc	gcagaaaaaa	aggatctcaa	3780
	gaagatcctt	tgatcttttc	tacgggggtct	gacgctcagt	ggaacgaaaa	ctcacgttaa	3840
	gggatttttg	tcatgagatt	atcaaaaagg	atcttcacct	agatcctttt	aaattaaaaa	3900
20	tgaagtttta	aatcaatcta	aagtatatat	gagtaaaact	ggtctgacag	ttaccaatgc	3960
	ttaatcagtg	aggcacctat	ctcagcgatc	tgtctatttc	gttcatccat	agttgcctga	4020
	ctccccgtcg	tgtagataac	tacgatacgg	gagggcttac	catctggccc	cagtgcctga	4080
	atgataccgc	gagaccacag	ctcaccggct	ccagatttat	cagcaataaa	ccagccagcc	4140
	ggaagggccg	agcgcagaag	tggtcctgca	actttatccg	cctccatcca	gtctattaat	4200
25	tggtgcccgg	aagctagagt	aagtagttcg	ccagttaata	gtttgcgcaa	cgttgttgcc	4260
	attgctacag	gcacgtgggt	gtcacgctcg	tcgtttggta	tggcttcatt	cagctccggg	4320
	tcccaacgat	caaggcgagt	tacatgatcc	cccatgttgt	gcaaaaaagc	ggttagctcc	4380
	ttcggctctc	cgaatctctt	cagaagtaag	ttggccgcag	tggtatcact	catgggtatg	4440
	gcagcactgc	ataattctct	tactgtcatg	ccatccgtaa	gatgcttttc	tgtgactggg	4500
30	gagtaactca	ccaagtcatt	ctgagaatag	tgtatgcggc	gaccgagttg	ctcttgcccg	4560
	gcgtcaaac	gggataatac	cgcgccacat	agcagaactt	taaaagtgc	catcattgga	4620
	aaacgttctt	cggggcgaaa	actctcaagg	atcttaccgc	tggtgagatc	cagttcgatg	4680
	taacccactc	gtgcacccaa	ctgatcttca	gcacctttta	ctttcaccag	cgtttctggg	4740
	tgagcaaaaa	caggaaggca	aaatgccgca	aaaaagggaa	taagggcgac	acggaaatgt	4800
35	tgaatactca	tactcttctc	ttttcaatat	tattgaagca	tttatcaggg	ttattgtctc	4860
	atgagcggat	acatatttga	atgtatttga	aaaaataaac	aaaagagttt	gtagaaacgc	4920
	aaaaaggcca	tccgtcagga	tggtccttctg	cttaatttga	tgcttgccag	tttatggcgg	4980
	gcgtcctgcc	cgccaccctc	cgggccggtg	cttcgcaacg	ttcaaatccg	ctcccgccgg	5040
	atgtgtctta	ctcaggagag	cgttcaccga	caaacaacag	ataaaacgaa	aggcccagtc	5100
40	tttcgactga	gcctttcgtt	ttatttgatg	cctggcagtt	ccctactctc	gcatggggag	5160
	acccacact	accatcggcg	ctacggcggt	tcacttctga	gttcggcatg	gggtcaggtg	5220
	ggaccacccg	gctactgccg	ccaggcaaat	tctgttttat	cagaccgctt	ctgcgttctg	5280
	attttaatctg	tatcaggctg	aaaatcttct	ctcatccgcc	aaaacagaag	ctagcggccg	5340
	atccccaaaa	aaaaaaaaaa	aaaaaaaaaa	aagagtcag	agtggccccc	ccgttccgcg	5400
45	ccgggggggg	gggggggggg	ggacactttc	ggacatctgg	tcgacctcca	gcacgggggg	5460
	aaaaaaaaaa	aacaaagttt	cgcccgaggt	actggctcgac	ctccgaagtt	gggggggg	5517

<210> 24

50 <211> 5699

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: pHL3236

<400> 24

```

5  cctctcataa tagacggcac agcctcatta agcccgggaa tgatgatggg tatgttcaac 60
   atgctgagta cagtgttggg agtctcaatc ctgaatcttg ggcaaaagag atacaccaa 120
   accacatact ggtgggatgg acttcagtc tctgatgatt ttgctctcat cgtgaatgca 180
   ccaaatcatg agggaataca agcgggagtg gatagattct acagaacctg caagctagtt 240
   gggatcaata tgagcaagaa aaagtcctat ataaatagga caggaacatt cgaattcaca 300
10 agctttttct accgctatgg atttgtagcc aatttttagta tggagttgcc cagctttgga 360
   gtatcaggaa ttaatgaatc ggctgatatg agcattggag taacagtgat aaagaataac 420
   atgataaaca atgatcttgg accggcaaca gcccaaatgg ctctccaatt attcatcaag 480
   gactacagat atacataccg gtgtcacagg ggagacacac aaatccaaac gaggaggtca 540
   ttcgagctaa agaagctgtg ggagcagacc cgctcaaagg caggactgtt ggtttcagat 600
15 ggcggaccaaa acctgtacaa cattcggaat ctccacatcc cggaagtgtt cctgaaatgg 660
   gaactaatgg atgaagacta tcagggaaga ctttghtaat ccatgaacct gtttgcagt 720
   cataaggaaa ttgaatctgt aaacaatgct gcggtaatgc cagcccatgg tccagccaaa 780
   agcatggaat atgatgctgt ggcaactaca cactcttggg tccctaagag aaaccgttcc 840
   attctcaata cgagtcaaag gggaatcctt gaggatgaac aaatgtacca gaagtgttgc 900
20 aacctattcg agaaattctt ccctagcagc tcatacagaa gaccagttgg aatttccagt 960
   atggtggagg ccatggtgtc tagggcccgg attgatgcac gaattgactt cgagtctgga 1020
   aggattaaga aggaagagtt tgctgagatc atgaagatcc cccgggctgc aggaattcga 1080
   tatcaagctt cgacgaattt ctgccattca tccgcttatt atcaattatt caggcgtagc 1140
   accaggcgtt taagggcacc aataactgcc ttaaaaaaat tacgccccgc cctgccactc 1200
25 atcgcagtac tgttgtaatt cattaagcat tctgccgaca tgggaagccat cacaaacggc 1260
   atgatgaacc tgaatcgcca gcggcatcag caccttgcg ccttgcgtat aatatttgcc 1320
   catggtgaaa acgggggcga agaagtgtgc catattggcc acgtttaaat caaaactggt 1380
   gaaactcacc cagggattgg ctgagacgaa aaacatattc tcaataaacc ctttagggaa 1440
   ataggccagg ttttcaccgt aacacgccac atcttgcgaa tatatgtgta gaaactgccg 1500
30 gaaatcgctg tggattcac tccagagcga tgaaaacgtt tcagtttgct catggaaaac 1560
   ggtgtaacaa ggggtgaacac tatcccatat caccagctca ccgtctttca ttgccatacg 1620
   gaattccgga tgagcattca tcaggcgggc aagaatgtga ataaaggccg gataaaactt 1680
   gtgcttattt ttctttacgg tctttaaaaa ggccgtaata tccagctgaa cggctctggt 1740
   ataggtacat tgagcaactg actgaaatgc ctcaaaatgt tctttacgat gccattggga 1800
35 tatatcaacg gtggtatatc cagtgtattt tttctccatt ttagcttctt tagctcctga 1860
   aaatctcgtc gaagcttatc gataccgtcg acctcgaggg ggggcccggg acggcctgca 1920
   aattttcaag aagatcattt ttcagctgat ctcgttatct agaccatgga gcttagtgat 1980
   ggtgatggtg atgggatccc ttgtatagtt catccatgcc atgtgtaate ccagcagctg 2040
   ttacaaaactc aagaaggacc atgtggtctc tcttttcgtt gggatctttc gaaagggcag 2100
40 attgtgtgga caggtaatgg ttgtctggta aaaggacagg gccatcgcca attggggtgt 2160
   tctgctggta gtggtcggcc agctgcacgc cgccgtctc gatgttgttg cgggtcttga 2220
   agttggcctt gattccgttc ttctgcttgt cagccatgat gtatacattg tgtgagttat 2280
   agttgtattc caatttgtgt ccaagaatgt ttccatcttc tttaaaatca atacctttta 2340
   actcgattct attaacaagg gtatcacctt caaacttgac ttcagcacgt gtctttagt 2400
45 tcccgctcatc tttgaaaaat atagttcttt cctgtacata accttcgggc atggcactct 2460
   tgaaaaagtc atgctgttct atatgatctg ggtatcttga aaagcattga acaccataag 2520
   tgaaaagtagt gacaagtgtt ggccatgga caggtagttt tccagtagtg caaataaatt 2580
   taagggtaa gtttccgtat gttgcatcac cttcacctc tccactgaca gaaaatttgt 2640
   gcccattaac atcaccatct aattcaacaa gaattgggac aactccagtg aaaagttctt 2700
50 ctcttttact catggtggac tccttcgcga gtcgaggggg ggcccgttac acgtacgcgc 2760
   tcgagaacgt accttcgcgc tcgcgatgtg gatgtcactc agtgagtgat tatctacctt 2820
   gtttctactc ccccccaact tcggaggtcg accagtactc cgggcgaaac tttgtttttt 2880
   ttttttcccc cgatgctgga ggtcgaccag atgtccgaaa gtgtccccc ccccccccc 2940

```

	ccccggcgcg	gaacggcggg	gccactctgg	actctttttt	tttttttttt	tttttttttt	3000
	ggggatcggc	cgctagcttc	tgttttggcg	gatgagagaa	gatttttcagc	ctgatacaga	3060
	ttaaatacaga	acgcagaagc	ggtctgataa	aacagaattt	gcctggcggc	agtagcgcg	3120
	tgggtcccacc	tgaccccatg	ccgaactcag	aagtgaaacg	ccgtagcgcc	gatggtagtg	3180
5	tggggtctcc	ccatgcgaga	gtagggaaact	gccaggcatc	aaataaaaacg	aaaggctcag	3240
	tcgaaagact	gggcctttcg	ttttatctgt	tgtttgctcg	tgaacgctct	cctgagtagg	3300
	acaaatccgc	cgggagcgga	tttgaacggt	gcgaagcaac	ggcccggagg	gtggcgggca	3360
	ggacgcccgc	cataaactgc	caggcatcaa	attaagcaga	aggccatcct	gacggatggc	3420
	ctttttgcgt	ttctacaaaac	tcttttggtt	atttttctaa	atacattcaa	atatgtatcc	3480
10	gctcatgaga	caataaccct	gataaatgct	tcaataatat	tgaaaaagga	agagtatgag	3540
	tattcaacat	ttcctgtctg	cccttattcc	cttttttgcg	gcattttgcc	ttcctgtttt	3600
	tgctcaccca	gaaacgctgg	tgaaagttaa	agatgctgaa	gatcagttgg	gtgcagcagt	3660
	gggttacatc	gaactggatc	tcaacagcgg	taagatcctt	gagagttttc	gccccgaaga	3720
	acgtttttcca	atgatgagca	ctttttaaagt	tctgctatgt	ggcgcgggat	tatcccgtgt	3780
15	tgacgcccgg	caagagcaac	tcggtcgccg	catacactat	tctcagaatg	acttggttga	3840
	gtactcacca	gtcacagaaa	agcatcttac	ggatggcatg	acagtaagag	aattatgcag	3900
	tgctgccata	accatgagtg	ataacactgc	ggccaactta	cttctgacaa	cgatcggagg	3960
	accgaaggag	ctaaccgctt	ttttgcacaa	catgggggat	catgtaactc	gccttgatcg	4020
	ttgggaaccg	gagctgaatg	aagccatacc	aaacgacgag	cgtgacacca	cgatgcctgt	4080
20	agcaatggca	acaacgttgc	gcaaaactatt	aactggcgaa	ctacttactc	tagcttcccg	4140
	gcaacaatta	atagactgga	tggaggcgga	taaagttgca	ggaccacttc	tgcgctcggc	4200
	ccttcgggct	ggctgggtta	ttgctgataa	atctggagcc	ggtgagcgtg	ggtctcgcgg	4260
	tatcattgca	gcactggggc	cagatggtaa	gccctcccgt	atcgtagtta	tctacacgac	4320
	ggggagtcag	gcaactatgg	atgaacgaaa	tagacagatc	gctgagatag	gtgcctcact	4380
25	gattaagcat	tggtaactgt	cagaccaagt	ttactcatat	atactttaga	ttgatttaaa	4440
	acttcatttt	taattttaaaa	ggatctaggt	gaagatcctt	tttgataatc	tcatgaccaa	4500
	aatcccttaa	cgtgagtttt	cgttccactg	agcgtcagac	cccgtagaaa	agatcaaagg	4560
	atcttcttga	gatccttttt	ttctgcgcgt	aatctgctgc	ttgcaaacaa	aaaaaccacc	4620
	gctaccagcg	gtggtttgtt	tgccggatca	agagctacca	actctttttc	cgaaggtaac	4680
30	tggcttcagc	agagcgcaga	taccaaatac	tgtccttcta	gtgtagccgt	agttaggcca	4740
	ccacttcaag	aactctgtag	caccgcctac	atacctcgct	ctgctaatac	tgttaccagt	4800
	ggctgctgcc	agtggcgata	agtcgtgtct	taccgggttg	gactcaagac	gatagttacc	4860
	ggataaggcg	cagcggtcgg	gctgaacggg	gggttcgtgc	acacagccca	gcttgagcgg	4920
	aacgacctac	accgaactga	gatacctaca	gcgtgagcta	tgagaaagcg	ccacgcttcc	4980
35	cgaagggaga	aaggcggaca	ggtatccggt	aagcggcagg	gtcggaaacag	gagagcgcac	5040
	gagggagctt	ccagggggaa	acgcctggta	tctttatagt	cctgtcgggt	ttcgccacct	5100
	ctgacttgag	cgtcgatttt	tgtgatgctc	gtcagggggg	cggagcctat	ggaaaaacgc	5160
	cagcaacgcg	gcctttttac	ggttcctggc	cttttgctgg	ccttttgctc	acatgttctt	5220
	tcttgctgta	tcccctgatt	cattaatgca	ggtcacgata	ctttctggcg	agtccccgtg	5280
40	cggagtcgga	gagcgcctcc	tgagcgcgcg	tgcggcccga	gaggtcgcgc	ctggccggcc	5340
	ttcgggtccct	cgtgtgtccc	ggtcgtagga	ggggccggcc	gaaaatgctt	ccggctcccc	5400
	ctctggagac	acgggcccgc	cccctgcgtg	tggcacgggc	ggccggggag	gcgtccccgg	5460
	cccggcgctg	ctcccgcgtg	tgtcctgggg	ttgaccagag	ggccccgggc	gctccgtgtg	5520
	tggctgcgat	ggtggcgctt	ttggggacag	gtgtccgtgt	cgcgcgtcgc	ctgggcccgc	5580
45	ggcgtggctg	gtgacgcgac	ctcccggccc	cgggggaggt	atatcttttcg	ctccgagtcg	5640
	gcattttggg	ccgcggggtt	attagtagaa	acaagggtat	ttttctttac	ctagctagg	5699

Claims

1. A recombinant influenza virus for high-yield expression of incorporated
5 foreign gene(s), which is genetically stable in the absence of any helper
virus and which comprises at least one viral RNA segment being a
bicistronic RNA molecule coding for two genes in tandem arrangement
(tandem RNA segment), in said tandem RNA segment one of the standard
viral genes being in covalent junction with a foreign, recombinant gene
10 and said tandem RNA segment having an upstream splice donor and a
downstream splice acceptor signal surrounding the proximal coding region.
2. The recombinant influenza virus of claim 1, wherein the tandem RNA
segment contains one of the standard viral genes in distal mRNA position
15 behind a foreign, recombinant gene in proximal position, or vice versa,
both in antisense orientation with regard to the viral RNA as present within
the virus.
3. The recombinant influenza virus of claim 1 or 2, wherein at least one
20 of the regular viral RNA segments is replaced by a tandem RNA segment,
preferably the replaced regular viral RNA segment is selected from the
neuraminidase segment, hemagglutinin segment and NS segment.
4. The recombinant influenza virus of claims 1 to 3, wherein the splice
25 donor and splice acceptor signals are selected from sequences as
present in influenza WSN segment 7 and 8 or other partially effective
splice reactin substrates.
5. The recombinant influenza virus of claim 4, wherein the splice donor
30 and splice acceptor signals are selected from sequences as present in
influenza WSN segment 7.

6. The recombinant influenza virus according to claims 1 to 5, wherein one or more of the regular viral RNA segments, differing from said at least one tandem RNA segment, comprises a vRNA encoding a foreign gene which may or may not be in covalent connection to one of the viral genes, and preferably one or more of the regular viral RNA segments has (have) been deleted and replaced by a tandem vRNA encoding in addition a foreign gene.

7. The recombinant influenza virus according to claims 1 to 6, in which the terminal viral RNA sequences of one or more of the regular segments and/or of the at least one tandem RNA segment, which are active as the promoter signal, have been modified by nucleotide substitutions in up to five positions, resulting in improved transcription rates of both the vRNA promoter as well as the cRNA promoter as present in the complementary sequence.

8. The recombinant influenza virus of claim 7, wherein the 12 nucleotide conserved influenza 3' terminal sequence has been modified by replacement of one to three nucleotides occurring in said sequence at positions 3, 5 and 8 relative to the 3' end by other nucleotides, and/or wherein the 13 nucleotide conserved influenza 5' terminal sequence has been modified by replacement of one or two nucleotides occurring in said sequence at positions 3 and 8 by other nucleotides.

9. The recombinant influenza virus of claim 8, wherein the replacements in the 3' terminal nucleotide sequence comprises the modifications G3A and C8U.

10. The recombinant influenza virus of claim 9, wherein the replacements in the 3' terminal nucleotide sequence comprises the modifications G3A, U5C and C8U, or G3C, U5C and C8G.

11. The recombinant influenza virus of claim 10, which comprises a 3' terminal nucleotide sequence of (5')-CCUGUUUCUACU-3'.
12. The recombinant influenza virus according to claims 7 to 12, wherein the 5' terminal nucleotide sequence comprises the modifications U3A and A8U resulting in a 5'-terminal sequence of 5'-AGAAGAAUCAAGG.
13. The recombinant influenza virus according to claims 1 to 12, which is a recombinant influenza A virus.
14. The recombinant influenza virus according to claims 1 to 13, in which the foreign gene(s) in the tandem RNA segment code for proteins and/or glycoproteins which are secreted from cells infected with the recombinant virus.
15. The recombinant influenza virus according to claims 1 to 13, in which the foreign gene(s) in the tandem RNA segment code for proteins or artificial polypeptides designed to support an efficient presentation of inherent epitopes at the surface of infected cells, for stimulation of a B cell and/or T cell response.
16. The recombinant influenza virus according to claims 1 to 13, in which the foreign gene(s) in the tandem RNA segment is a nucleotide sequence causing viral attenuation.
17. The recombinant influenza virus of claim 16, wherein the foreign gene is coding for part of or for the entire viral neuraminidase gene in antisense orientation.
18. The recombinant influenza virus of claim 17, wherein the neuraminidase gene in antisense orientation is attached to the hemagglutinin vRNA

segment, and optionally another gene or reporter gene is encoded in a second tandem vRNA, preferably in conjunction with NS2.

19.A method for the production of recombinant influenza viruses as
5 defined in claims 1 to 18 comprising

(a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, in antisense or in sense tandem design,

(b) followed by infection with an influenza carrier strain constructed to include flanking ribozyme target sequences in the corresponding viral RNA
10 segment, and

(c) thereafter selective vRNA inactivation through ribozyme cleavage.

20.A pharmaceutical composition comprising a recombinant influenza virus according to claims 1 to 18, preferably a recombinant influenza virus
15 of claims 16 to 18.

21.Use of a recombinant influenza virus according to claims 1 to 18, preferably a recombinant influenza virus of claims 16 to 18, for preparing a medicament for vaccination purposes.

20

22.The use according to claim 21, wherein the medicament

(a) is suitable against influenza and/or against other infections;

(b) is present in form of inactivated preparations; and/or

(c) is present in form of live recombinant viruses.

25

23.Use of a recombinant influenza virus according to claims 1 to 18 for preparing agents for somatic gene therapy.

24. Use of a recombinant influenza virus according to claims 1 to 18 for
30 preparing agents, for transfer and expression of foreign genes into cells infected by such viruses.

25. Use of a recombinant influenza virus according to claims 1 to 18 for preparing agents for transfer and expression of RNA molecules into cells infected by such viruses.

- 5 26. The use of claim 24, wherein the RNA molecules to be expressed are antisense sequences or double-strand sequences relative to the target cell cellular mRNA molecules, and/or the agent is suitable for sequence-specific gene silencing, preferably by antisense RNA or RNA interference mechanisms.

10

27. The use according to claims 23 to 26, wherein the agents are applicable in *ex vivo* and *in vivo* application schemes.

- 15 28. A method for the production of proteins or glycoproteins which comprises utilizing a recombinant influenza virus according to claims 1 to 19 as expression vector.

29. The method of claim 28, wherein the production is performed in cell culture cells or in fertilized chicken eggs.

20

30. A method for preventing and/or treating influenza which comprises administering an effective amount of a recombinant influenza virus according to claims 1 to 18, preferably of a recombinant influenza virus according to claims 16 to 18, to the mammal to be treated.

25

31. A method for somatic gene therapy, which method comprises subjecting the organism to be treated with a recombinant influenza virus according to claims 1 to 18.

- 30 32. A method for transfer and expression of foreign genes into cells, and for transfer and expression of RNA molecules into cells, which method

comprises infecting the cells with a recombinant influenza virus according to claims 1 to 18.

33. Use of a recombinant influenza virus according to claims 1 to 18 for
5 preparing agents for immunotherapy, preferably for autologous immunotherapy.

34. A method for an immunotherapy which comprises *ex vivo* infection of immune cells, preferably dendritic cells, with a recombinant influenza virus
10 according to claims 1 to 18, and introduction of the transduced cells into the patient.

35. A method for the induction of antibodies which comprises utilizing a recombinant influenza virus according to claims 1 to 18 as an immunogen.
15

FIG. 1




FIG. 2






FIG. 3


pHL3196



pHL 3235



pHL 3224



pHL3236




FIG. 4



A**B**

FIG. 5




FIG. 6




FIG. 7




FIG. 8

